



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

and

$$\theta = (2\lambda - 1) \frac{\pi}{2} + \frac{q}{p} (4\mu \pm 1) \frac{\pi}{2}, \quad (\lambda = 1, 2, 3, \dots, \frac{q-1}{2}; \mu = 0, 1, 2, \dots, p-1);$$

determine the same set of points on the curve

$$\rho = a \cos \frac{p}{q} \theta,$$

where p and q are two odd integers without a common factor, and a is any constant.

471. Proposed by C. N. SCHMALL, New York City.

In the ellipse $x^2/a^2 + y^2/b^2 = 1$, an *equilateral* hexagon is inscribed with two sides parallel to the major axis. In the major auxiliary circle the same thing is done. If H_1 and H_2 be the sides of the hexagons, and e the eccentricity of the ellipse, show that $H_1 : H_2 :: 4 - 2e^2 : 4 - e^2$

CALCULUS.

390. Proposed by WILSON L. MISER, University of Minnesota.

Show that the triangle whose area is a constant and whose perimeter is a minimum is equilateral.

391. Proposed by H. B. PHILLIPS, Massachusetts Institute of Technology.

If $0 < \lambda < 1$ and $0 < x < \pi$, show that the function $(\sin \lambda x)/(\sin x)$ increases as x increases.

392. Proposed by HORACE OLSON, Student at The University of Chicago.

Two right circular cylinders of radii a and b respectively, are placed so that their axes intersect at right angles. Find the volume common to them.

MECHANICS.

314. Proposed by C. N. SCHMALL, New York City.

A rectangular box of height h , and having a plane mirror for its bottom, contains a quantity of water of unknown height x . In the lid are two small apertures distant $2a$ from each other. A ray of light entering one aperture with an angle of incidence ϕ , emerges, after refraction and reflection, through the other aperture. If μ be the index of refraction of water, show that the height of the water is

$$x = \frac{(h \tan \phi - a)}{\left[\tan \phi - \frac{\sin \phi}{(\mu^2 - \sin^2 \phi)^{\frac{1}{2}}} \right]}.$$

SOLUTIONS OF PROBLEMS.

ALGEBRA.

427. Proposed by CLIFFORD N. MILLS, Brookings, S. Dak.

If $r \sin (\theta + \alpha) = m$, and $r \cos (\theta + \beta) = n$, show that

$$r = \frac{\sqrt{m^2 + n^2 - 2mn \sin (\alpha - \beta)}}{\cos (\alpha - \beta)}.$$

SOLUTION BY J. H. KELLOGG, Oberlin College.

From trigonometry, using only the positive square root, we know that

$$\begin{aligned}
\sin (A+B) &= \sqrt{(\sin A \cos B + \cos A \sin B)^2} \\
&= \sqrt{\sin^2 A (1 - \sin^2 B) + (1 - \sin^2 A) \sin^2 B + 2 \sin A \cos A \sin B \cos B} \\
&= \sqrt{\sin^2 A + \sin^2 B + 2 \sin A \sin B (\cos A \cos B - \sin A \sin B)} \\
&= \sqrt{\sin^2 A + \sin^2 B + 2 \sin A \sin B \cos (A+B)}.
\end{aligned}$$

From the given equations we have

$$\sin (\theta + \alpha) = \frac{m}{r} \quad \text{and} \quad \sin \left[\frac{\pi}{2} - (\theta + \beta) \right] = \frac{n}{r}.$$

Hence letting

$$A = (\theta + \alpha) \quad \text{and} \quad B = \left[\frac{\pi}{2} - (\theta + \beta) \right],$$

we have

$$(A+B) = \left[\frac{\pi}{2} + (\alpha - \beta) \right].$$

Substituting, we have

$$\begin{aligned}
&\sin \left[\frac{\pi}{2} + (\alpha - \beta) \right] \\
&= \sqrt{\sin^2 (\theta + \alpha) + \sin^2 \left[\frac{\pi}{2} - (\theta + \beta) \right] + 2 \sin (\theta + \alpha) \sin \left[\frac{\pi}{2} - (\theta + \beta) \right] \cos \left[\frac{\pi}{2} + (\alpha - \beta) \right]},
\end{aligned}$$

or

$$\cos (\alpha - \beta) = \sqrt{\frac{m^2}{r^2} + \frac{n^2}{r^2} - 2 \frac{m}{r} \frac{n}{r} \sin (\alpha - \beta)}.$$

Solving for r , we have

$$r = \frac{\sqrt{m^2 + n^2 - 2mn \sin (\alpha - \beta)}}{\cos (\alpha - \beta)}.$$

Also solved by B. J. BROWN, HERBERT N. CARLTON, NATHAN ALTSHILLER, C. E. HORNE, C. N. SCHMALL, PAUL CAPRON, J. A. CAPARO, FRANK IRWIN, FRANK R. MORRIS, S. A. JOFFE, V. M. SPUNAR, L. G. WELD, GEORGE W. HARTWELL, ELIJAH SWIFT, A. W. SMITH, JOSEPH B. REYNOLDS, RICHARD MORRIS, WALTER C. EELLS, ALBERT N. NAUER, ELMER SCHUYLER, and A. M. HARDING.

428. Proposed by FRANK IRWIN, University of California.

If the roots of the equation

$$x^n - na_1x^{n-1} + \binom{n}{2}a_2x^{n-2} + \cdots = 0$$

are all real, the condition that they should all be equal is $a_1^2 = a_2$. A proof of the sufficiency of the condition is readily obtained from a consideration of derivatives. A proof is desired not based on such considerations.

SOLUTION BY ELIJAH SWIFT, University of Vermont.

We are to show that if the roots are all real, and if $a_1^2 = a_2$, the roots are all equal. For that purpose write the above equation in the form